

A systematic review and meta-analysis of acupuncture in *in vitro* fertilisation

T El-Toukhy,^a SK Sunkara,^a M Khairy,^a R Dyer,^a Y Khalaf,^a A Coomarasamy^{a,b}

^a Assisted Conception Unit, Guy's and St Thomas' Hospital, London, UK ^b Department of Public Health and Epidemiology, University of Birmingham, Birmingham, UK

Correspondence: Dr A Coomarasamy, Assisted Conception Unit, Guy's Hospital, London SE1 9RT, UK. Email: arricoomar@blueyonder.co.uk

Accepted 5 June 2008. Published OnlineEarly 23 July 2008.

Background Numerous randomised studies have reported pregnancy outcome in women who received acupuncture during their *in vitro* fertilisation (IVF) treatment cycle.

Objective The objective of this study was to conduct a systematic review with meta-analysis of the trials of acupuncture during IVF treatment on the outcomes of clinical pregnancy and live birth rates.

Search strategy Searches were conducted in MEDLINE, EMBASE, Cochrane Library, ISI Proceedings and SCISEARCH.

Selection criteria All randomised controlled trials that evaluated the effects of acupuncture compared with no treatment or sham acupuncture in women undergoing IVF–intracytoplasmic sperm injection treatment were included.

Data collection and analysis Study selection, quality appraisal and data extraction were performed independently and in duplicate. A sensitivity analysis was conducted where the meta-analysis was restricted to trials in which sham acupuncture was used in the control group. Meta-regression analysis was used to explore the association between study characteristics and pregnancy rates.

Main results Thirteen relevant trials, including a total of 2500 women randomised to either acupuncture or control group, were identified. No evidence of publication bias was found (Begg's

test, $P = 0.50$). Five trials ($n = 877$) evaluated IVF outcome when acupuncture was performed around the time of transvaginal oocyte retrieval, while eight trials ($n = 1623$) reported IVF outcome when acupuncture was performed around the time of embryo transfer (ET). Meta-analysis of the five studies of acupuncture around the time of egg collection did not show a significant difference in clinical pregnancy (relative risks [RR] = 1.06, 95% CI 0.82–1.37, $P = 0.65$). Meta-analysis of the eight studies of acupuncture around the time of ET showed no difference in the clinical pregnancy rate (RR = 1.23, 95% CI 0.96–1.58, $P = 0.1$). Live birth data were available from five of the eight studies of acupuncture around the time of ET. Meta-analysis of these studies did not show a significant increase in live birth rate with acupuncture (RR = 1.34, 95% CI 0.85–2.11). Using meta-regression, no significant association between any of the studied covariates and clinical pregnancy rate was found ($P > 0.05$ for all covariates).

Conclusion Currently available literature does not provide sufficient evidence that adjuvant acupuncture improves IVF clinical pregnancy rate.

Keywords Acupuncture, IVF, oocyte retrieval, embryo transfer, clinical pregnancy, randomised trials.

Please cite this paper as: El-Toukhy T, Sunkara S, Khairy M, Dyer R, Khalaf Y, Coomarasamy A. A systematic review and meta-analysis of acupuncture on *in vitro* fertilisation. BJOG 2008;115:1203–1213.

Introduction

Approximately three-quarters of all *in vitro* fertilisation (IVF) cycles fail.¹ Clinicians are constantly searching for ways to improve IVF results by methods other than replacing more embryos. Over the past decade, a number of treatment strategies have been designed to increase IVF success rate, primarily through improving the quality of embryos replaced or uterine environment or both. Many of these strategies, however, have been introduced into clinical practice and

promoted to patients before a clear evidence of benefit has been established.^{2,3}

One of the treatments that has gained popularity in recent years is the use of alternative therapies, and in particular acupuncture, as an adjunctive treatment to improve IVF outcome.^{4,5} The scientific rationale for using acupuncture during IVF treatment has not been fully accepted but focuses mainly on its potential role in enhancement of uterine receptivity through increased blood flow⁶ and quiescence.⁷ In addition, a perceived reduction in anxiety and stress after

acupuncture^{8–10} has often been cited as an additional justification for its use.

Numerous studies have already reported IVF outcome in women who had received acupuncture during their IVF treatment cycle. These studies had variable design and have generally yielded inconclusive or conflicting results, rendering the clinical decision whether to recommend or omit the use of acupuncture during IVF difficult to make.^{11–13} A recent systematic review examined the role of acupuncture at the time of embryo transfer (ET);¹⁴ it excluded numerous randomised studies that evaluated the role of acupuncture at the time of egg retrievals.^{15–19} Furthermore, new randomised evidence²⁰ has emerged since the publication of this review, casting uncertainty on the conclusions of the review.

In the present study, we sought to conduct a systematic review of randomised trials involving the use of acupuncture during IVF treatment, to generate a more precise estimate of the effect of acupuncture on IVF outcome.

Materials and methods

Literature search

We searched MEDLINE (1966 to January 2008), EMBASE (1974 to January 2008), Cochrane Library (2007:4) and SCISEARCH (1974 to January 2008) for relevant studies. A combination of Medical Subject Headings (MeSH) and text words was used to generate two subsets of citations, one including studies of acupuncture ('acupuncture', 'acupressure', 'moxibustion', 'electroacupuncture', 'auricular-acupuncture' and 'acupunc*') and the other studies of IVF and intracytoplasmic sperm injection (ICSI) ('in vitro fertilization', 'Fertilization-in-vitro', 'intracytoplasmic-sperm-injection', 'sperm-injections-intracytoplasmic', 'assisted reproductive techniques', 'embryo transfer' and 'embryo implantation'). These subsets were combined using 'AND' to generate a subset of citations relevant to our research question. We also searched ISI Proceedings for conference abstracts, and ISRCTN Register and Meta-register for RCTs (*mRCT*) for continuing and archived randomised controlled trials. The reference lists of relevant primary and review articles were examined to identify cited articles not captured by electronic searches. Articles frequently cited were used in the Science Citation Index to identify additional citations. Authors were contacted to obtain missing information. No language restrictions was placed in any of our searches.

Study selection

Studies were selected if the target population was women undergoing IVF with or without ICSI treatment, the therapeutic intervention was any accepted regimen of (needle or laser) acupuncture compared with no or sham (placebo) acupuncture and pregnancy outcome was reported from a single IVF cycle per woman randomised to receive either

acupuncture or control intervention. Studies with a crossover design were excluded.

The outcome measures of interest were clinical pregnancy and live birth rates per IVF cycle started. For the purpose of this review, clinical pregnancy was defined as the ultrasound identification of an intrauterine gestational sac after IVF treatment.

Studies were selected in a two-stage process. First, the titles and abstracts from the electronic searches were scrutinised by two reviewers independently (T.E.-T. and S.K.S.), and full manuscripts of all citations that were likely to meet the pre-defined selection criteria were obtained. Second, final inclusion or exclusion decisions were made on examination of the full manuscripts. In cases of duplicate publication, the most recent and complete versions were selected. The assessment of English language manuscripts was performed independently by two reviewers (T.E.-T. and S.K.S.) and other language manuscripts by people who had command of the language. Any disagreements about inclusion were resolved by consensus or arbitration by a third reviewer (A.C.).

The selected studies were assessed for methodological quality using the components of study design that are related to internal validity.^{21,22} Information on the adequacy of randomisation, concealment of allocation, blinding, the use of sham (or placebo) acupuncture and intention-to-treat analysis was sought by examining the full-text articles and by contacting the authors if clarification was needed.

For our review, we accepted any standard method of delivering sham acupuncture, including (a) superficial needling of the true acupuncture points, (b) application of true acupuncture in the wrong location or in points designed for *other* medical conditions, (c) use of blunt (placebo) needles or (s) use of sham laser acupuncture.

Data extraction and statistical analysis

Study characteristics such as population features and interventions (e.g. exact regimen of acupuncture, time of commencement and duration of treatment) were extracted from each study. Outcome data from each study were extracted in 2×2 structured tables using an intention-to-treat approach, and the results were pooled and expressed as relative risks (RR) with 95% CI. Heterogeneity of treatment effects was evaluated graphically using forest plot and statistically using chi-square test.

We proceeded to perform meta-analyses separately for the two broad groups of studies defined by the timing of delivery of acupuncture, that is around the time of transvaginal oocyte retrieval (TVOR) and around the time of ET. We also attempted to do a sensitivity analyses based on whether sham acupuncture was used in the control group. For our meta-analysis, we used a random effects model because of the encountered heterogeneity of the trials' characteristics and populations studied.²³ Meta-regression was then used to

explore the possible sources of the observed heterogeneity between studies.²⁴

To assess for publication bias, we performed a funnel plot analysis using Begg's test.²⁵ As the meta-analysis did not involve subjecting patients to an intervention, and data were extracted from pre-existing literature, there was no need for obtaining approval by our local research ethics committee. All statistical analyses were performed using Stata 8.0 (StatCorp LP, TX, USA) and RevMan 4.2.10 (Cochrane Collaboration, Oxford, UK) softwares.

Results

The literature search yielded 83 citations, of which 43 were selected for retrieval. Figure 1 summarises the process of literature identification and selection. Of the 43 full manuscripts examined, 13 articles,^{15–20,26–32} including a total of 2500 women, met the selection criteria for our review. The methodological quality of the included trials is summarised in Table 1. No evidence of publication bias or related biases was suggested from the funnel plot analysis (Begg's test, $P = 0.50$).

Five of the 13 included trials ($n = 877$) provided pregnancy outcome data when acupuncture was performed around the time of TVOR,^{15–19} while the remaining eight trials ($n = 1623$)

evaluated pregnancy outcome when acupuncture was performed around the time of ET.^{20,26–32} Each group of trials was considered separately.

Acupuncture at the time of TVOR

Main study characteristics

Five trials ($n = 877$) reported IVF outcome when acupuncture was performed at the time of TVOR. Tables 2 and 3 show the features of the five trials and the acupuncture points used, respectively. The mean age of participants ranged from 30.5 to 34.4 years. All five studies were performed in Europe: three were conducted in Sweden,^{15,16,18} one was conducted in Denmark¹⁷ and one was conducted in Austria.¹⁹ Three trials were performed in a single centre,^{17–19} while the remaining two were multicentre trials.^{15,16}

Four of the five trials^{15,17–19} were designed to assess the pain-relieving effects of acupuncture used at the time of TVOR compared with conventional analgesia. Only one study used¹⁶ the IVF pregnancy rate as the primary outcome. Three of the five trials^{17–19} were powered to detect a clinically significant difference in pain intensity or wellbeing after TVOR between the study groups, and in one study, *a priori* power calculation was not described.¹⁵ The only study¹⁶ that was powered to detect a 10% difference in the clinical pregnancy rate between the study groups was terminated prematurely




Figure 1. Study selection process for the systematic review of acupuncture in women undergoing IVF-ICSI treatment.

Table 1. Quality of studies included in the systematic review of acupuncture use during IVF

Study	Method of randomisation	Allocation concealment	Blinding	Placebo intervention	ITT	Comparability at baseline
Stener-Victorin <i>et al.</i> ¹⁵ (1999)	Not mentioned	Adequate	No	No	Yes	Unclear
Paulus <i>et al.</i> ²⁶ (2002)	Computerised randomisation	Adequate	Single blind	No	Yes	Yes
Stener-Victorin <i>et al.</i> ¹⁶ (2003)	Not mentioned	Adequate	No	No	No	Unclear
Paulus <i>et al.</i> ^{27,33} (2003)	Not mentioned	Adequate	No	Yes	Yes	Unclear
Humaidan and Stener-Victorin ¹⁷ (2004)	Not mentioned	Adequate	No	No	Yes	Yes
Gejervall <i>et al.</i> ¹⁸ (2005)	Computerised randomisation	Unclear	No	No	No	Unclear
Dieterle <i>et al.</i> ³⁰ (2006)	Not mentioned	Adequate	Double blind	Yes	Yes	Yes
Westergaard <i>et al.</i> ²⁸ (2006)	Unclear	Adequate	No	No	No	Yes
Smith <i>et al.</i> ²⁹ (2006)	Block randomisation	Adequate	Single blind	Yes	Yes	Yes
Sator-Katzenschlager <i>et al.</i> ¹⁹ (2006)	Computerised randomisation	Unclear	Double blind	Yes	Yes	Yes
Benson <i>et al.</i> ³² (2006)	Not mentioned	Unclear	No (except laser groups)	No (except laser group)	Yes	Yes
Domar <i>et al.</i> ³¹ (2006)	Not mentioned	Adequate	Single blind	No	Yes	Yes
Craig <i>et al.</i> ²⁰ (2007)	Computerised randomisation	Yes	Single blind	No	No	Yes

ITT, intention-to-treat analysis.

based on the results of an interim analysis, which showed no difference in the clinical pregnancy rate between the two groups. None of the five studies used a sham acupuncture technique in the control group. One study¹⁸ described the IVF protocol used. The quality of the embryos replaced and day of ET were reported in only one of the five studies,¹⁹ although the day of ET was not standardised among the study participants in that study.

IVF treatment outcome

For the clinical pregnancy rate, data were available from all five trials. Using the random effects model, pooling of the results from all five trials showed no significant difference in the clinical pregnancy rate between the acupuncture and the controls groups (RR = 1.06, 95% CI 0.82–1.37, $P = 0.65$; Figure 2).

Acupuncture around the time of ET

Main study characteristics

Eight trials ($n = 1623$) compared IVF outcome when acupuncture was performed around the time of ET with that in a control group. Tables 4 and 5 show features of these trials and the acupuncture points used, respectively. Four of the eight studies^{20,27,31,32} were published as conference abstracts only, while the remaining four were published as full reports.

Four of the eight studies were conducted in Europe (three in Germany^{26,27,30} and one in Denmark,²⁸ three in USA^{20,31,32} and one in Australia²⁹).

Seven of the eight trials were performed as single-centre trials,^{26–32} while one study was a multicentre trial.²⁰ All studies were designed as two-arm trials, except the study of Westergaard *et al.*,²⁸ which included three arms (two intervention groups and one control group), and that of Benson *et al.*,³² which included five arms (two intervention and three control groups). Unlike the studies examining the effect of acupuncture at the time of TVOR, all eight trials were designed with the objective to assess the effect of acupuncture performed at the time of ET on IVF outcome. Three studies^{26,28,30} described the IVF protocol used. The quality of the embryos replaced and day of ET were reported in only three^{26,28,29} and five studies,^{20,26,28,30,32} respectively. One of the eight studies²⁷ did not mention the number of embryos replaced in the study groups.

Choice of therapeutic intervention

All eight studies used traditional needle acupuncture, and none used electroacupuncture. In addition, the study of Benson *et al.*³² used laser acupuncture in one of the five arms of the trial (half of those randomised to the intervention group). Six of the eight studies used a needle acupuncture technique similar to that described by Paulus *et al.*,²⁶ in which the treatment group received the acupuncture treatment for 25 minutes before and 25 minutes after ET. In addition, women in the treatment group received a third acupuncture session on day 9 of ovarian stimulation in the study of Smith *et al.*²⁹ or 2 days after ET in the study Westergaard *et al.*²⁸ In

Table 3. The acupuncture points used in the studies of acupuncture performed at the time of TVOR

Author	Acupuncture points								
	LI 4	TE 5	ST 29	GV 20	ST 36	SP 6	KI 11	LI 10	AA
Stener-Victorin <i>et al.</i> ¹⁵ (1999)	Yes	Yes	Yes	Yes	Yes	—	—	—	—
Stener-Victorin <i>et al.</i> ¹⁶ (2003)	Yes	Yes	Yes	Yes	Yes	—	—	—	—
Humaidan and Stener-Victorin ¹⁷ (2004)	Yes	—	—	Yes	—	Yes	—	—	—
Gejervall <i>et al.</i> ¹⁸ (2005)	Yes	—	Yes	Yes	Yes	—	Yes	Yes	—
Sator-Katzenschlager <i>et al.</i> ¹⁹ (2006)	—	—	—	—	—	—	—	—	Yes

AA, auricular acupuncture; GV, governor vessel; KI, kidney; LI, large intestine; SP, spleen; ST, stomach; TE, triple energiser.

CI 0.99–1.76, $P = 0.06$; Figure 5). Exclusion of the study of Dieterle *et al.*³⁰ did not change the meta-analysis result (RR = 1.18, 95% CI 0.88–1.59, $P = 0.15$).

Exploration of sources of heterogeneity between studies of acupuncture around the time of ET. We attempted to explore effect of the potential sources of observed heterogeneity between the studies on pregnancy outcome using meta-regression analysis. We included in the meta-regression model study characteristics that were considered potentially significant, namely allocation concealment, nature of acupuncture technique used, who administered the acupuncture treatment, number of acupuncture sessions delivered to each woman in the intervention groups and use of sham acupuncture in the control groups. No significant association between any of these covariates and clinical pregnancy rate was found ($P > 0.05$), although the analysis was limited by the small number of studies included.

Discussion

Complimentary and alternative therapies are widely used, with acupuncture ranking among the most popular therapies being used.^{5,34,35} As a result, a link between acupuncture and

IVF outcome is likely to be of considerable interest to clinicians and patients alike.

Advocates of acupuncture have suggested that it could improve IVF outcome through a number of possible mechanisms, including a central sympathoinhibitory effect, resulting in increased uterine blood flow, which in turn might improve endometrial receptivity;⁶ stimulation of beta-endorphins release, which could influence steroid hormone secretion;^{36–39} and a direct, or endocrine-mediated, inhibitory effect on uterine activity.⁴¹

This systematic review and meta-analysis used the clinical pregnancy and live birth rates as indicators of the effect of acupuncture performed during IVF treatment on cycle outcome. The findings of our review fail to show a significant improvement in the clinical pregnancy or live birth rates associated with the use of acupuncture whether performed at the time of TVOR or around the time of ET. According to our results, the true effect of acupuncture performed at the time of TVOR on IVF outcome ranges from up to 13% relative reduction to a 24% relative increase in the chance of a clinical pregnancy and that of acupuncture performed around the time of ET ranges from up to 4% relative reduction to a 58% relative increase in the chance of a clinical pregnancy per IVF cycle started compared with no acupuncture.




Figure 2. Meta-analysis of the studies evaluating the effect of acupuncture administered around the time of TVOR on the clinical pregnancy rate in women undergoing IVF.




Figure 4. Meta-analysis of the studies evaluating the effect of acupuncture administered around the time of ET on the clinical pregnancy rate in women undergoing IVF when no sham acupuncture technique was used in the control groups.

undergoing IVF and found that acupuncture treatment did not inhibit uterine activity as previously suggested.

Another suggested benefit from acupuncture, which might potentially lead to improvement in IVF success rate, was reduction of stress levels and improvement in psychological wellbeing in women undergoing IVF.^{28,30,45} Interestingly, the only two randomised trials that attempted to test this hypothesis failed to provide supportive evidence. The study of Smith *et al.*²⁹ found more women in the control group reporting sense of ‘relaxation’ and feeling ‘calm and peaceful’ after ET (67 and 64%, respectively) compared with the acupuncture group (51 and 55%, respectively). Furthermore, Domar *et al.*³¹ reported no significant differences between the study and the control groups in optimism levels after ET.

The choice of the control intervention also varied between the studies that examined the effect of acupuncture performed around the time of ET and could have contributed to the conflicting results reported in these studies. Paulus *et al.*²⁷ and Myers¹² raised the possibility that acupuncture might exaggerate pregnancy rates after IVF through a placebo effect. Contrary to this suggestion, pooling the results of the four studies in which no placebo intervention was employed in the control group yielded an effect size closer to the line of

unity than the studies that employed a sham acupuncture technique. Furthermore, different forms of sham acupuncture were employed in the four studies that examined the effect of acupuncture at the time of ET on IVF outcome. The lack of a reproducible and reliable sham acupuncture technique that does not affect the acupoints (e.g. by acupressure or *shiatsu*) and is devoid of any negative effect¹² undermines the reliability of the results of these studies, may explain to a certain extent the significant degree of heterogeneity present among these studies and underlines one of the many difficulties faced in conducting such trials.^{42,46,47}

Given the cost, relative invasiveness of acupuncture, potential for harm and the significant variation in the inherent features of the published studies, women embarking on IVF should be advised that based on current knowledge, there is insufficient evidence that receiving acupuncture during IVF treatment (whether at time of oocyte collection or ET) improves cycle outcome. Our review shows clearly that despite the publication of 13 trials of acupuncture during IVF, well-designed and conducted research into the efficacy and cost-effectiveness of acupuncture carried out as an adjunct to IVF treatment is still needed before clinicians could recommend its use.




Figure 5. Meta-analysis of the studies evaluating the effect of acupuncture administered around the time of ET on the clinical pregnancy rate in women undergoing IVF when sham acupuncture technique was used in the control groups.

Disclosure of interests

The authors state that they have no conflict of interest relating to any pharmaceutical, clinical, consumer or other groups.

Contribution to authorship

A.C. conceived the review; T.E.-T., S.K.S. and M.K. performed literature searches, study selection and data extraction. T.E.-T. and A.C. performed the analysis and wrote the initial draft. M.K., S.K.S., R.D. and Y.K. critically revised the manuscript.

Acknowledgement

We acknowledge Prof Craig and Prof Domar for providing additional study data. ■

References

- 1 HFEA Code of Practice, 7th edn. London: HFEA, 2007.
- 2 Urman B, Yakin K, Balaban B. Recurrent implantation failure in assisted reproduction: how to counsel and manage. A. General considerations and treatment options that may benefit the couple. *Reprod Biomed Online* 2005;11:371–81.
- 3 De Sutter P. Rational diagnosis and treatment in infertility. *Best Pract Res Clin Obstet Gynaecol* 2006;20:647–64.
- 4 Hayhoe S, Box H. A questionnaire on medical acupuncture practice. *Acupunct Med* 1997;15:96–9.
- 5 Thomas K, Nicholl J, Coleman P. Use and expenditure on complementary medicine in England: a population based survey. *Complement Ther Med* 2001;9:2–11.
- 6 Stener-Victorin E, Waldenström U, Andersson SA, Wikland M. Reduction of blood flow impedance in uterine arteries of infertile women with electro-acupuncture. *Hum Reprod* 1996;11:1314–17.
- 7 Kim J, Shin K, Na C. Effect of acupuncture treatment on uterine motility and cyclooxygenase-2 expression in pregnant rates. *Gynecol Obstet Invest* 2000;50:225–30.
- 8 Domar A, Seibel M, Benson H. The mind/body program for infertility: a new behavioral treatment approach for women with infertility. *Fertil Steril* 1990;53:246–9.
- 9 Dong J. Research on the reduction of anxiety and depression with acupuncture. *Am J Acupunct* 1993;21:327–30.
- 10 MacPherson H, Thomas K. Short term reactions to acupuncture: a cross-sectional survey of patient reports. *Acupunct Med* 2005;23:112–20.
- 11 White A. A review of controlled trials of acupuncture for women's reproductive health care. *J Fam Plann Reprod Health Care* 2003;29:233–6.
- 12 Myers ER. Acupuncture as adjunctive therapy in assisted reproduction: remaining uncertainties. *Fertil Steril* 2006;85:1362–3.
- 13 Collins J. The play of chance. *Fertil Steril* 2006;85:1364–7.
- 14 Manheimer E, Zhang G, Udoff L, Haramati A, Langenberg P, Berman BM, *et al.* Effects of acupuncture on rates of pregnancy and live birth among women undergoing in vitro fertilization: systematic review and meta-analysis. *BMJ* 2008;336:545–9.
- 15 Stener-Victorin E, Waldenström U, Nilsson L, Wikland M, Olof Janson P. A prospective randomised study of electro-acupuncture versus alfentanil as anaesthesia during oocyte aspiration in in-vitro fertilization. *Hum Reprod* 1999;14:2480–4.
- 16 Stener-Victorin E, Waldenström U, Wikland M, Nilsson L, Hägglund L, Lundeberg T. Electro-acupuncture as a perioperative analgesic method and its effects on implantation rate and neuropeptide Y concentrations in follicular fluid. *Hum Reprod* 2003;18:1454–60.
- 17 Humaidan P, Stener-Victorin E. Pain relief during oocyte retrieval with a new short duration electro-acupuncture technique—an alternative to conventional analgesic methods. *Hum Reprod* 2004;19:1367–72.
- 18 Gejervall AL, Stener-Victorin E, Möller A, Olof Janson P, Werner C, Bergh C. Electro-acupuncture versus conventional analgesia: a comparison of pain levels during oocyte aspiration and patients' experiences of well-being after surgery. *Hum Reprod* 2005;20:728–35.
- 19 Sator-Katzenschlager SM, Wöfler MM, Kozek-Langenecker SA, Sator K, Sator PG, Li B, *et al.* Auricular electro-acupuncture as an additional perioperative analgesic method during oocyte aspiration in IVF treatment. *Hum Reprod* 2006;21:2114–20.
- 20 Craig L, Criniti A, Hansen K, Marshall L, Soules M. Acupuncture lowers pregnancy rates when performed before and after embryo transfer. *Fertil Steril* 2007;88 (Suppl 1):S40.
- 21 Undertaking Systematic Reviews of Research on Effectiveness. CRD's Guidance for Carrying Out or Commissioning Reviews. CRD Report Number 4, 2nd edn. York, UK: NHS Centre for Reviews and Dissemination, University of York, 2001.
- 22 Juni P, Altman DG, Egger M. Systematic reviews in health care: assessing the quality of controlled clinical trials. *BMJ* 2001;323:42–6.
- 23 Ades AE, Lu G, Higgins JPT. The interpretation of random-effects meta-analysis in decision models. *Med Decis Making* 2005;25:646–54.
- 24 Thompson S, Higgins J. How should meta-regression analyses be undertaken and interpreted? *Stat Med* 2002;21:1559–73.
- 25 Hayashino Y, Noguchi Y, Fukui T. Systematic evaluation and comparison of statistical tests for publication bias. *J Epidemiol* 2005;15:235–43.
- 26 Paulus WE, Zhang M, Strehler E, El-Danasouri I, Sterzik K. Influence of acupuncture on the pregnancy rate inpatients who undergo assisted reproduction therapy. *Fertil Steril* 2002;77:721–4.
- 27 Paulus W, Zhang M, Strehler E, Sterzik K. Motility of the endometrium after acupuncture treatment. *Fertil Steril* 2003;80 (Suppl 3):131.
- 28 Westergaard LG, Mao Q, Kroglund M, Sandrini S, Lenz S, Grinsted J. Acupuncture on the day of embryo transfer significantly improves the reproductive outcome in infertile women: a prospective, randomised trial. *Fertil Steril* 2006;85:1341–6.
- 29 Smith C, Coyle M, Norman RJ. Influence of acupuncture stimulation on pregnancy rates for women undergoing embryo transfer. *Fertil Steril* 2006;85:1352–8.
- 30 Dieterle S, Ying G, Hatzmann W, Neuer A. Effect of acupuncture on the outcome of in vitro fertilization and intracytoplasmic sperm injection: a randomised, prospective, controlled clinical study. *Fertil Steril* 2006;85:1347–51.
- 31 Domar A, Meshay I, Kelliher J, Wang S, Alper M. The impact of acupuncture on IVF outcome. *Fertil Steril* 2006;86 (Suppl 2):S378–9.
- 32 Benson M, ElKind-Hirsch K, Theall A, Fong K, Hogan R, Scott R. Impact of acupuncture before and after embryo transfer on the outcome of invitro fertilization cycles: a prospective single blind randomised study. *Fertil Steril* 2006;86 (Suppl 2):S135.
- 33 Paulus W, Zhang M, Strehler E, Seybold B, Sterzik K. Placebo-controlled trial of acupuncture effects in assisted reproductive therapy. *Hum Reprod* 2003;18 (Suppl 1):xviii18–xviii19.
- 34 Eisenberg D, Davis R, Ettner S, Appel S, Wilkey S, Van Rompay M, *et al.* Trends in alternative medicine use in the United States 1990–1997: results of a follow-up national survey. *JAMA* 1998;280:1569–75.
- 35 Ernst E, White A. The BBC survey of complimentary medicine use in the United Kingdom. *Complement Ther Med* 2000;8:32–6.

- 36 Aso T, Motohashi T, Murata M, Nishimura T, Kakizaki K. The influence of acupuncture stimulation on plasma levels of LH, FSH, progesterone and estradiol in normally ovulating women. *Am J Chin Med* 1976;4:391–401.
- 37 Han J, Terenius L. Neurochemical basis of acupuncture analgesia. *Annu Rev Pharmacol Toxicol* 1982;22:193–220.
- 38 Takeshige C, Oka K, Mizuno T, Hisamitsu T, Luo CP, Kobori M, *et al*. The acupuncture point and its connecting central pathway for producing acupuncture analgesia. *Brain Res Bull* 1993;30:53–67.
- 39 Petti F, Bangrazi A, Liguori A, Reale G, Ippoliti F. Effects of acupuncture on immune response related to opioid-like peptides. *J Tradit Chin Med* 1998;18:55–63.
- 40 Chang R, Chung PH, Rosenwaks Z. Role of acupuncture in the treatment of female infertility. *Fertil Steril* 2002;78:1149–53.
- 41 Schultz K, Chalmers I, Hayes R, Altman D. Empirical evidence of bias: dimensions of methodological quality associated with estimates of treatment effects in controlled trials. *JAMA* 1995;273:408–12.
- 42 Anderson BJ, Rosenthal L. Acupuncture and IVF controversies. *Fertil Steril* 2006;87:1000.
- 43 Wang W, Check JH, Liss JR, Choe JK. A matched controlled study to evaluate the efficacy of acupuncture for improving pregnancy rates following in vitro fertilization-embryo transfer. *Clin Exp Obstet Gynecol* 2007;34:137–8.
- 44 Stener-Victorin E, Humaidan P. Use of acupuncture in female infertility and a summary of recent acupuncture studies related to embryo transfer. *Acupunct Med* 2006;24:157–63.
- 45 Luo H, Meng F, Jia Y, Zhao X. Clinical research on the therapeutic effect of the electroacupuncture treatment in patients with depression. *Psychiatry Clin Neurosci* 1998;52:338–40.
- 46 Park J, White A, Stevinson C, Ernst E, James M. Validating a new non-penetrating sham acupuncture device: two randomised controlled trials. *Acupunct Med* 2002;20:168–74.
- 47 Dincer F, Linde K. Sham interventions in randomised clinical trials of acupuncture—a review. *Complement Ther Med* 2003;11:235–42.